

60th Annual Scientific Session & Expo

E276

JACC April 5, 2011

Volume 57, Issue 17



CARDIAC FUNCTION AND HEART FAILURE

HIGH SENSITIVITY C-REACTIVE PROTEIN AS AN INDEPENDENT PREDICTOR OF PROGRESSIVE MYOCARDIAL FUNCTIONAL DETERIORATION: THE LONGITUDINAL MULTI-ETHNIC STUDY OF ATHEROSCLEROSIS

ACC Poster Contributions

Ernest N. Morial Convention Center, Hall F

Sunday, April 03, 2011, 3:30 p.m.-4:45 p.m.

Session Title: Biomarkers and Prognostication in Heart Failure

Abstract Category: 24. Myocardial Function/Heart Failure—Clinical Nonpharmacological Treatment

Session-Poster Board Number: 1054-7

Authors: Eui-Young Choi, Raymond Yan, Veronica Fernandes, Anders Opdahl, Antoinette Gomes, Andre Almeida, Colin Wu, Kiang Liu, Jeffrey J. Carr, Robyn L. McClelland, David A. Bluemke, Joao AC Lima, Johns Hopkins Hospital, Baltimore, MD

Objectives: In this study, we used longitudinal MRI follow-up analysis to examine whether C-reactive protein (CRP) levels relate to progressive myocardial functional deterioration as a potential mechanism of incident heart failure.

Background: Systemic inflammation has been linked to the development of heart failure in population studies including MESA (Multi-Ethnic Study of Atherosclerosis) but little evidence exists regarding potential mechanism of this relationship. Method: Regional myocardial functional data from MESA participants who had baseline CRP measurement and also underwent tagged cardiac MRI both at baseline and at five-year follow-up (n=785) were analyzed. Left ventricular (LV) midwall peak systolic circumferential strain (Ecc), of which a more negative value denotes stronger regional myocardial function, was measured, while Ecc change was calculated as the difference between baseline and follow-up Ecc.

Results: During the follow-up period, participants with elevated CRP experienced a decrease in strain, independent of age, gender and ethnicity (B=0.08; Δ Ecc change per 1mg/L CRP change, 95% CI 0.04-0.13, p<0.001, Model 1), and additionally beyond systolic blood pressure, diabetes, smoking status, body mass index, current medication and glomerular filtration rate (B=0.10, 95% CI 0.05-0.14, p<0.001, Model 2). The relationship remained statistically significant after further adjustment for LV mass, coronary calcium score and interim clinical coronary events (B=0.10, 0.05-0.15, p<0.001).

Conclusions: Higher CRP levels are related to progressive myocardial functional deterioration independent of subclinical atherosclerosis and clinical coronary events in asymptomatic individuals without previous history of heart disease.